

Kinetic Model for Batch Cellulase Production by *Aspergillus terreus* at Different Levels of Dissolved Oxygen Tension using Oil Palm Empty Fruit Bunch Fibre as Substrate

Mahdi Shahriarinnour^{1*} Mohd Noor Abdul Wahab²

Abstract: In this study, the effect of different levels of dissolved oxygen tension (d.o.t) on the production of three main components of extracellular cellulases (FPase, CMCase and β -glucosidase) at a fixed agitation speed by *Aspergillus terreus* was investigated. The results from the model based on logistic and Luedeking-Piret equations fit well with the experimental data, confirming that the models were appropriate for describing and representing growth and cellulase production at various dissolved oxygen tension (d.o.t) levels. The models showed that the production of FPase and CMCase were growth associated processes. Cell growth and cellulase production were approximately two-fold higher in a stirred tank bioreactor compared with a shake-flask culture. At a d.o.t of 55% air saturation, cell growth and cellulase production were higher than at low d.o.t (40% air saturation) or high d.o.t (80% air saturation). The highest activities of FPase (2.33 U ml⁻¹), CMCase (51.10 U ml⁻¹), and β -glucosidase (16.18 U ml⁻¹) were obtained at a d.o.t of 55% air saturation, yielding overall productivities of 19.40, 425.00, and 67.40 U l.h⁻¹, respectively.

Keywords—Kinetics; Cellulase; *Aspergillus terreus*; Oil palm empty fruit bunch; Stirred tank bioreactor

I. INTRODUCTION

Cellulase production is a major factor in the hydrolysis of cellulosic material and it is essential to unlocking the economic potential of biomass resources. *Aspergillus* species play a key role in decay and decomposition due to their ability to produce a wide range of enzymes. Although cellulase production has been reported for several *Aspergillus* species [1], only a few studies have reported on the production of cellulase from *Aspergillus terreus* [2].

1 Mahdi shahriarinnour is with the Department of Microbiology, Science and Research Branch, Islamic Azad University, Guilan, Iran (corresponding author's phone:+989359310560; fax:+981317233987 ; e-mail: Mahdi.shahriari@srbiau.ac.ir).

2 Department of Bioprocess Technology, Faculty of Biotechnology and Biomolecular Sciences, University of Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia (e-mail: mmoorwa@biotech.upm.edu.my).

The amount of cellulase production is affected by the composition of the nutrient medium and environmental conditions, which can be inducers or repressors of cell density and growth rate [3]. In addition, cellulase production is influenced by the hydrodynamic conditions in the bioreactor [4]. The effect of d.o.t in the culture during fermentation (at a fixed agitation speed) on the synthesis of cellulase has not been adequately studied. Oil palm empty fruit bunch (OPEFB) fiber is an appropriate renewable raw material for bioconversion into value added products because it is easily accessible, plentiful and rich in lignocellulose. Improved cellulase production by *A. terreus* has been previously reported and the optimized conditions in shake-flask and a stirred tank bioreactor were achieved through response surface methodology [5], [6]. The kinetic study is useful in obtaining kinetic parameter values that describes the kinetic behavior. Hence, the main objective of this study was to investigate the kinetics of *A. terreus* growth and cellulase production, which are FPase, CMCase, and β -glucosidase using simplified model in a stirred tank bioreactor using oil palm empty fruit bunch (OPEFB) fiber. The experimental data from batch fermentation of cellulase at various d.o.t levels on the growth and the production of the three major components of cellulase [6] were analyzed in order to form the basis for a kinetic model of the process.

II. MATERIALS AND METHODS

A. Microorganism

The mould *Aspergillus terreus* used in this study for cellulase production was isolated from a sample collected from the oil palm empty fruit bunch (OPEFB) compost [7]. It was grown on potato dextrose agar (PDA) at 30 °C for 7 days to allow the development of spores and then stored at 4 °C until use in inoculum preparation.

B. Medium and Inoculum Preparation

The basal medium as proposed by Mandels and Weber [8] was used for cellulase production. In

all experiments, yeast extract (8 g L⁻¹) was added as the only nitrogen source and delignified OPEFB fibre (13.90 g L⁻¹) as a carbon source to the basal medium for cellulase production [6]. The OPEFB fibres obtained from an oil palm processing factory (Sri Ulu Langat Palm, Dengkil, Selangor, Malaysia) were delignified by physico-chemical pretreatment by soaking in phosphoric acid and then exposed to hydrothermal treatment at 160 °C for 10 min, followed by biological treatment with effective microorganisms [9]. The spore suspension containing, on average 6 × 10⁷ spores mL⁻¹, was used as an inoculum in all fermentation.

C. Crude Enzyme Extraction

After an appropriate time of incubation, the cultures were harvested at 24 h intervals by centrifugation at 15,000 rpm at 4 °C for 15 min using a refrigerated ultracentrifuge (SORVALL RT7 PLUS). The supernatant was then analyzed for soluble protein and extracellular enzyme activities.

D. Fermentations in Bioreactor

The fermentations were carried out in 2 L stirred tank bioreactor (Biostat B, Melsungen, Germany) and the vessel was made of 5 mm thick borosilicate glass. The vessel had an internal diameter of 0.13 m and liquid height of 0.12 m. Two six-bladed Rushton turbine impellers with a diameter (D) of 52 mm were mounted on the impeller shaft used for agitation. The bioreactor was equipped with temperature and dissolved oxygen controllers. Air was supplied into the culture via a single air sparger (0.1 mm internal diameter). During the fermentation, agitation speed (N) was fixed at 225 rev/min (impeller tip speed = πND = 0.613 m/s), and d.o.t in the culture broth was controlled using a sequential cascade control of airflow rate. The maximum and minimum set points of permitted airflow rates were 1.5 L/min and 0.1 L/min, respectively. A polarographic dissolved oxygen probe (Ingold, Urdorf, Switzerland) was used to measure d.o.t in the culture. In all cases, d.o.t was effectively controlled to within ± 2 % of the required set points (40, 55, 60, and 80 % saturation). The initial pH (5.5) of the culture were adjusted to appropriate values either by addition of 1 N HCl or 1N NaOH. The temperature within the bioreactor was controlled at 29 °C. Spore suspension (15 mL) was inoculated into the bioreactor containing 1.5 L medium. During the fermentation, samples were withdrawn at regular time intervals for analysis.

E. Analytical Procedure

To estimate mycelium concentration, the chemical method based on the measurement of glucosamine was selected while the physical

separation of mycelium from the OPEFB fibres for measurement was not possible [10]. Carboxymethylcellulase (CMCase) activity was determined by measuring spectrophotometrically the reducing sugar produced from 2% (w/v) carboxymethyl-cellulose, while filter-paper-hydrolysing (FPase) activity was determined by estimating the reducing sugar liberated from filter paper [11]. Meanwhile, β-glucosidase was determined using the method described by Wood and Bhat [11]. In this method, *p*-nitrophenol released from *p*-nitrophenyl-β-D-glucopyranoside (Fluka) was measured using a spectrophotometer (Shimadzu, UV-1601 PC). One unit of β-glucosidase activity is defined as 1 μmol *p*-nitrophenol liberated/mL of enzyme/min, while the specific activity is defined as units/mg protein. Protein content was determined by the method of Bradford using bovine serum albumin as a standard [12].

F. Mathematical Approach

The following simplified batch fermentation kinetic models for cell growth and product formation based on logistic and Luedeking-Piret equations were used to evaluate the kinetics of production of the three major components of cellulase (FPase, CMCase, and β-glucosidase) by *A. terreus*,

$$\text{Cell growth} \quad \frac{dx}{dt} = \left[\mu_{\max} \left(1 - \frac{x}{x_{\max}} \right) \right] x \quad (1)$$

$$\text{CMCase} \quad \frac{dA}{dt} = p \left(\frac{dx}{dt} \right) + qx \quad (2)$$

$$\text{FPase} \quad \frac{dB}{dt} = a \left(\frac{dx}{dt} \right) + bx \quad (3)$$

$$\beta\text{-glucosidase} \quad \frac{dC}{dt} = e \left(\frac{dx}{dt} \right) + fx \quad (4)$$

where x is the cell concentration (g L⁻¹); x_{\max} is the maximum cell concentration (g L⁻¹); μ_{\max} is the maximum specific growth rate (h⁻¹); A is the CMCase activity (U mL⁻¹); B is the FPase activity (U mL⁻¹); C is β-glucosidase activity (U mL⁻¹); p is the growth associated coefficient for CMCase formation (U g-cell⁻¹); q is the non-growth associated coefficient for CMCase formation (U g-cell.h⁻¹); a is the growth associated coefficient for FPase formation (U g-cell⁻¹); b is the non-growth associated coefficient for FPase formation (U g cell.h⁻¹); e is the growth associated coefficient for β-glucosidase formation (U g-cell⁻¹); and f is the non-growth associated coefficient for β-glucosidase formation (U g-cell.h⁻¹).

The kinetic models (Eq. 1-4) were fitted to the experimental data using a non-linear regression with a Marquadt algorithm using MATLAB computer software (MATLAB 7.2, The MathWorks, 2006 U.S.).

III.RESULTS AND DISCUSSION

A.Distribution of Cellulase in *A. terreus*

The distribution of the three major components of cellulase (FPase, CMCase, and β -glucosidase) of *A. terreus* was identified in fermentation using soluble CMC as carbon source. Table 1 shows the maximum production and location of FPase, CMCase and β -glucosidase in *A. terreus* using soluble CMC as the carbon source. The FPase and CMCase of *A. terreus* were fully extracellular, while β -glucosidase was both an extracellular (cell free) and a cell bound (surface bound and intracellular) enzyme. The amount of extracellular β -glucosidase of *T. reesei* is less than in the mycelium at all stages of growth. Unlike *T. reesei*, the distribution of β -glucosidase in *A. terreus* was dependent on the age of the mycelium [3]. Most β -glucosidase was intracellular when the culture was in the active stage of growth [3]. It is possible that this enzyme is released by autolysis, i.e., when the culture is in the stationary phase.

TABLE I
Localization of Cellulase in *Aspergillus terreus* Grown in Shake-Flask Culture

Enzyme	Intracellular or cell bound (U ml ⁻¹)	Extracellular or free (U ml ⁻¹)
FPase	0	0.71 ±0.03
CMCase	0	8.16 ±0.71
B-glycosidase	10.23±0.74	5.85 ±0.53

B. Production of Cellulase in a Shake-Flask

Typical time course results for free cellulase production by *A. terreus* in a shake-flask cultures containing the basal medium as proposed by Mandels and Weber [8] was used with an addition of yeast extract (8 g L⁻¹) as the only nitrogen source and treated OPEFB fibres (13.90 g L⁻¹) as a carbon source are shown in Fig. 1.




Fig. 1 Comparison between the calculated and experimental data for shake-flask fermentation of cellulase by *A. terreus*. \diamond , Cell concentration; \bullet , FPase activity; \blacktriangle , CMCase

activity; \square , β -glucosidase activity; (—) data calculated according to Eq. 1-4

TABLE II
The Kinetics Parameter Values of Fermentation in Shake-Flask and Bioreactor for Production of All Three Main Components of Cellulase Production by *A. terreus*

Kinetic Parameter	Bioreactor at different d.o.t levels (% saturation)				
	Shake-flask	40	55	60	80
X_{max} (g/l)	5.20 ± 0.02 ^a	11.54 ± 0.01 ^b	13.07 ± 0.03 ^c	12.68 ± 0.02 ^d	7.15 ± 0.01 ^e
μ_{max} (h ⁻¹)	0.0041 ± 0.0003 ^a	0.0069 ± 0.0002 ^b	0.0071 ± 0.0002 ^b	0.0070 ± 0.0001 ^b	0.0054 ± 0.0003 ^c
FPase	1.13 ± 0.01 ^a	1.95 ± 0.01 ^b	2.33 ± 0.04 ^c	2.24 ± 0.02 ^d	1.36 ± 0.03 ^e
B_{max} (U/ml)	0.238 ± 0.05 ^a	0.207 ± 0.03 ^b	0.287 ± 0.06 ^c	0.273 ± 0.07 ^d	0.161 ± 0.02 ^e
a (U/g cell)	0	0	0	0	0
b (U/g cell.h)	14.25 ± 0.04 ^a	45.6 ± 0.07 ^b	51.10 ± 0.12 ^c	49.2 ± 0.09 ^d	22.4 ± 0.06 ^e
CMCase	2.76 ± 0.05 ^a	3.55 ± 0.02 ^b	4.28 ± 0.05 ^c	3.83 ± 0.02 ^d	2.96 ± 0.03 ^e
p (U/g cell)	0	0	0	0	0
q (U/g cell.h)	9.86 ± 0.03 ^a	14.66 ± 0.05 ^b	16.18 ± 0.06 ^c	15.83 ± 0.04 ^d	11.5 ± 0.09 ^e
β -glucosidase	0.819 ± 0.06 ^a	0.515 ± 0.01 ^b	1.039 ± 0.08 ^c	0.856 ± 0.02 ^d	0.267 ± 0.04 ^e
C_{max} (U/ml)	0.118 ± 0.009 ^a	0.0049 ± 0.0004 ^b	0.0053 ± 0.0002 ^b	0.0051 ± 0.0006 ^b	0.0048 ± 0.0003 ^b
e (U/g cell)	0.217 ± 0.01 ^a	0.169 ± 0.02 ^b	0.178 ± 0.01 ^b	0.176 ± 0.03 ^b	0.19 ± 0.02 ^c
f (U/g cell.h)	2.74 ± 0.04 ^a	3.96 ± 0.02 ^b	3.90 ± 0.03 ^c	3.88 ± 0.02 ^c	3.13 ± 0.05 ^d
B_{max}/X_{max} (×1000 U/g)	1.89 ± 0.03 ^a	1.27 ± 0.01 ^b	1.23 ± 0.02 ^c	1.24 ± 0.01 ^c	1.60 ± 0.04 ^d
C_{max}/X_{max} (×1000 U/g)	178.2 ± 1.5 ^a	307.5 ± 1.9 ^b	367.5 ± 2.3 ^c	353.3 ± 3.2 ^d	214.5 ± 2.1 ^e
Yield (U FPase / g cellulose l.h)	9.41 ± 0.16 ^a	15.25 ± 0.22 ^b	19.41 ± 0.25 ^c	18.66 ± 0.23 ^d	11.33 ± 0.19 ^e
Productivity (U FPase / l.h)					

^{a-e} Values in the same row with different superscripts are significantly different ($P < 0.05$)

Values are the means of triplicate ± standard deviation

The CMCase and FPase activities were increased throughout the growth cycle and reached their maximum during the stationary phase. The rate of free β -glucosidase production was low during the initial stages of the fermentation and very rapid towards the end of the fermentation. In contrast to CMCase and FPase production, free β -glucosidase was produced during the stationary phase. This result suggests that the proposed models based on logistic and Luedeking-Piret equations are sufficient to simulate growth of *A. terreus* and the production of the cellulase complex.

The kinetics parameter values calculated by the models for fermentation in a shake-flask can be used to describe the kinetics of the production of all three main components of cellulase by *A. terreus* (Table 2). The non-growth associated rate constants for FPase (b) and CMCCase (q) are zero, suggesting that production of these enzymes was growth associated processes. Instead, the production of free β -glucosidase was a mixed process indicated by non-zero values of the growth associated (e) and non-growth associated rate constants (f).

C. Cellulase Production in the Bioreactor at Various Dissolved Oxygen Tension Levels

Growth and cellulase production of three major components of cellulase in the fermenter at different d.o.t levels by *A. terreus* are shown in Fig. 2. The models of growth and cellulase production in the stirred tank bioreactor were comparable to that achieved in a shake-flask. However, growth and cellulase production by *A. terreus* were significantly affected by the d.o.t levels during the fermentation. Growth and cellulase production were high at a d.o.t of 55% air saturation, while the growth was inhibited at high d.o.t (80% air saturation). The effect of d.o.t on cellulase production was similar to the effect on growth, indicating that cellulase production was directly associated to the rate of growth and cell concentration in the culture. Figure 2 also shows how the calculated points from the proposed models (Eq. 1-4) fit the experimental data that determine fermentation at various d.o.t levels. In all cases, the calculated data fit well with the experimental data with greater than 85 % confidence. Table 2 shows the evaluation of the function and the kinetic factor values of cellulase production in the stirred tank fermenter at different d.o.t levels. In all cases, the values of b and q are zero while the value of f is not zero. These results explain a previous outcome of the experiment using a shake-flask: the FPase and CMCCase production were growth associated processes, while the production of free β -glucosidase was a mixed-growth associated process. The highest x_{\max} , A_{\max} , B_{\max} and C_{\max} values were obtained at d.o.t levels of 55 % air saturation. In addition, the μ_{\max} was also higher at d.o.t levels of 55% air saturation. However, the values of a , p , e and f were significantly different at various d.o.t levels. During the fermentation, x_{\max} was different with various d.o.t levels in the culture. Therefore, the maximum amount of cellulase components should be determined per unit cell weight (B_{\max}/x_{\max} , A_{\max}/x_{\max} and C_{\max}/x_{\max}) to signify whether the fungi produced the enzyme efficiently in certain culture conditions or if the production was simply proportional to cell mass.




Fig. 2 Comparison between the calculated data and the experimental data for batch fermentation of cellulase by *A. terreus* in the bioreactor at different d.o.t levels. (a) Cell concentration; (b) FPase activity; (c) CMCCase activity; (d) β -glucosidase activity. ■ 40%, ▲ 55%, ◇ 60%, ○ 80%; (—) data calculated according to Eq.1-4

The values of B_{\max}/x_{\max} , A_{\max}/x_{\max} were not significantly different at various d.o.t levels (Table 2). This result suggests that CMCase and FPase production corresponded to cell weight. However, the value of C_{\max}/x_{\max} (1.60×1000 U/g) was the highest at 80% air saturation, while the value was not significantly different at 40%, 55% and 60% air saturation, signifying that the efficiency of cell to produce β -glucosidase was improved at very high d.o.t. The highest yields of FPase (367.50 U/g-cellulose and 19.41 U/l.h, respectively) were obtained at a d.o.t of 55% air saturation. Higher shear rates in the bioreactor disrupt the fungal mycelium and reduce cellulase production [13].

During excess oxygen conditions (80% air saturation), the production of CMCase and FPase were significantly reduced. Additionally, improved cellulase production at a d.o.t level that was higher than the critical value of dissolved oxygen (C_{crit}) was detected in *T. reesei* [14]. The ability of *A. terreus* to produce β -glucosidase (as measured by C_{\max}/x_{\max}) was increased at high d.o.t (80 % air saturation); however, growth was significantly inhibited, which resulted in a reduction in the maximum activity of free β -glucosidase. Typically, higher d.o.t and agitation intensity were required to enhance β -glucosidase production. The production of all three main components of cellulase in the stirred tank bioreactor, where the d.o.t was controlled at 55 % air saturation, was approximately two-fold higher than the production in a shake-flask. The high shear rate created by the agitation using an impeller may have caused excessive breakdown of the OPEFB fibers, which reduced the fiber size and increased the degree of attack by microbes and enzymes. In a shake-flask culture, the fungus grew in the form of a pellet with an atypical diameter between 0.5 and 1 mm, whereas in the bioreactor, the fungus grew in the form of mycelia. The low shear rate in a shake-flask culture facilitated growth into the form of a pellet, and this had been detected in other cellulase producers, such as *T. reesei* [13]. The pelletization of hyphae in the shake-flask fermentation led to severe mass transfer limitation, which was responsible for lower μ_{\max} compared with the μ_{\max} for growth in the bioreactor at 55% air saturation. However, the pelletization enhanced the leakage of enzyme from inside the pellet. Therefore, the reduction in cellulase production in the shake-flask culture was attributed to the partial lack of dissolved oxygen and to the leakage of enzymes from the internal parts of the pellet.

In the bioreactor, where a high d.o.t level (80% air saturation) was maintained, growth of *A. terreus* was inhibited. The influence of high d.o.t levels in the inhibition of microbial growth in the

bioreactor has been described by indicating that the surplus oxygen is, at some level, toxic to the growth of some microorganisms [15]. Superoxide radicals (O_2^-), which are critical to cell metabolism, are major inhibitors of cell growth [16]. A simple model using logistic and Luedeking-Piret equations was found sufficient to describe growth of *A. terreus* and the production of all three main components of cellulase. From the model, it was recognized that the production of FPase and CMCase were growth-associated, while free β -glucosidase production was mixed. In order to attain a high amount of β -glucosidase in the culture filtrate, the fermentation should be extended after growth has reached stationary phase. The cellulase produced by *A. terreus* exhibited a high level of β -glucosidase activity. The ratio of β -glucosidase to FPase for cellulase obtained in fermentation at 55% d.o.t was about 8:1, and this was significantly higher than the ratios for *T. reesei* [17], *T. lignorum* [18], *Fusarium oxysporum* [19], *Gliocladium virens* [20], and *Chaetomium globosum* [3], which were in the range 0.44 to 7.

IV. CONCLUSIONS

A considerable agreement among the calculated data and the experimental data for both cell growth and the production of all the three main components of cellulase was observed, suggesting that the recommended model based on logistic and Luedeking-Piret equations was sufficient to describe the growth of *A. terreus* and the three main components of cellulase production. FPase and CMCase production were growth-associated processes (fully extracellular) while the production of free β -glucosidase was a mixed-growth-associated process (both an extracellular and a cell bound). Whilst, the cell growth and cellulase production were higher at 55% saturation of d.o.t as compared to the production at low condition oxygen (40% saturation) and at very high d.o.t (80% saturation). The production cellulase in stirred tank bioreactor with d.o.t maintained at 55% air saturation was improved approximately two-fold compared to the production of the shake-flask culture.

ACKNOWLEDGEMENTS

The first author would like to extend his gratitude for the financial support generously provided by Malaysia's Ministry of Science, Technology and Innovation (MOSTI) under the project 02-01-04-SF0735.

REFERENCES

- [1] R. A. Lockington, L. Rodbourn, S. Barnett, C. J. Carter, and J. M. Kelly , “ Regulation by carbon and nitrogen sources of a family of cellulases in *Aspergillus nidulans*,” *Fungal Genetics and Biology*., vol. 37, no. 2, pp. 190-196, 2002.
- [2] G. Emtiazi, N. Naghavi, and A. Bordbar. “Biodegradation of lignocellulosic waste by *Aspergillus terreus*,” *Biodegradation*., vol.12, no. 4, pp. 259-263, 2001.
- [3] M. S. Umikalsom, A. B. Ariff, M. A. Hassan, M. A., and M. I. A. Karim , “Kinetics of cellulase production by *Chaetomium globosum* at different levels of dissolved oxygen tension using oil palm empty fruit bunch fibre as substrate,” *World Journal of Microbiology and Biotechnology*., vol. 14, no.4, pp. 491-498, 1998.
- [4] J. Garcia-Soto Mariano, E. Botello-Alvarez, H. Jimenez-Islas, J. Navarrete-Bolanos, E. Barajas-Conde, and R. Rico-Martínez, R , “Growth morphology and hydrodynamics of filamentous fungi in submerged cultures,” *Advances in Agricultural and Food Biotechnology*., vol. 2, no. 1, pp.17-34, 2006.
- [5] M. Shahriarinnour, M. N. A. Wahab, A. Arbakariya, M. Rosfarizan, and M. Shuhaimi, “Optimization of cellulase production by *Aspergillus terreus* under submerged fermentation using response surface methodology,” *Australian Journal of Basic and Applied Sciences*., vol. 4, no. 12, pp. 6101-6124, 2010.
- [6] M. Shahriarinnour, R. N. Ramanan, R. N., M. N. A. Wahab, M. Rosfarizan, M. Shuhaimi, M., and B. A. Arbakariya, , “Improved cellulase production by *Aspergillus terreus* using oil palm empty fruit bunch fibre as substrate in a stirred tank bioreactor through optimization of the fermentation conditions,” *BioResources* ., vol. 6, no. 3, pp. 2663-2675, 2011.
- [7] M. Shahriarinnour, M. N. A. Wahab, A. Arbakariya, M. Rosfarizan, M , “Screening, isolation and selection of cellulolytic fungi from oil palm empty fruit bunch fibre,” *Biotechnology*., vol. 10, no. 1, pp. 108-113, 2011.
- [8] M. Mandels, and J. Weber, J , “The production of cellulases,” *Advances in Chemistry Series*., vol.95, pp. 391-414, 1969.
- [9] M. Shahriarinnour, M. N. A. Wahab, M. Shuhaimi, M. Rosfarizan, and B. A. Arbakariya, , “Effect of various pretreatments of oil palm empty fruit bunch fibres for subsequent use as substrate on the performance of cellulase production by *Aspergillus terreus*,” *BioResources*., vol. 6, no. 1, pp. 291-307, 2011.
- [10] S. R. Khan, and R. N. Strange, “Evidence for the role of a fungal stimulant as a determinant of differential susceptibility of jute cultivars to *Colletotrichum corchori*,” *Physiological Plant Pathology*., vol. 5, no. 2, pp. 157-162, 1975.
- [11] T. M. Wood, and K. M. Bhat, “Methods for measuring cellulase activities,” *Methods in Enzymology*., vol. 160, no. 2, PP. 87-112, 1998.
- [12] M. M. Bradford, “A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein dye binding,” *Analytical Biochemistry*., vol. 72, no.1-2, pp. 248-254, 1976.
- [13] R. Lejeune, and G. V. Baron, , “Effect of agitation on growth and enzyme production of *Trichoderma reesei* in batch fermentation,” *Applied Microbiology and Biotechnology*., vol. 43, no. 2, pp. 249-258, 1995.
- [14] S. Lantz, F. Goedegebuur, R. Hommes, T. Kaper, B. Kelemen, C. Mitchinson, L. Wallace, J. Stahlberg, and E. Larenas, E, “*Hypocrea jecorina* CEL 6 A protein engineering,” *Biotechnology for Biofuels*., vol. 3, no. 1, pp. 1-13, 2010.
- [15] C. Lee, S. Lee, K. Jung, S. Katoh, and E. Lee, “High dissolved oxygen tension enhances heterologous protein expression by recombinant *Pichia pastoris*,” *Process Biochemistry*., vol. 38, no. 8, pp.1147-1154, 2003.
- [16] B. Lund, T. Baird-Parker, and G. Gould, “*The Microbiological Safety and Quality of Food*,” Springer US, 2000.
- [17] R. Doppelbauer, H. Esterbauer, W. Steiner, R. M. Lafferty, and H. Steinmuller, “The use of lignocellulosic wastes for production of cellulase by *Trichoderma reesei*,” *Applied Microbiology and Biotechnology*., vol. 26, no. 5, pp. 485-494, 1987.
- [18] M. Baig, “Cellulolytic enzymes of *Trichoderma lignorum* produced on banana agro-waste: Optimisation of culture medium and conditions,” *Journal of Scientific and Industrial research*., vol. 64, no. 1, pp. 57-60, 2005.
- [19] G. Ramanathan, S. Banupriya, and D. Abirami, D , “Production and optimization of cellulase from *Fusarium oxysporum* by submerged fermentation,” *Journal of Scientific and Industrial Research*., vol. 69, no.6, pp. 454-459, 2010.
- [20] J. Gomes, I. Gomes, H. Esterbauer, W. Kreiner, and W. Steiner, W, “Production of cellulases by a wild strain of *Gliocladium virens*: Optimization of the fermentation medium and partial characterization of the enzymes,” *Applied Microbiology and Biotechnology*., vol. 31, no. 5-6, pp. 601-608, 1989.